

Aunt Cathy's Guide to:

# Thinking About OTHER Nutrition Issues in Diabetes



**AUNT CATHY**

Cathy Breedon PhD, RD, CSP, FADA  
Prenatal/Pediatric Nutrition Specialist  
Clinical Nutrition Specialist  
MeritCare Medical Center, Fargo, ND  
and UND School of Medicine

## Introduction:

Blood sugar control is the key focus of nutrition and diet planning for people with diabetes. Historically the most attention has been paid to adjusting the intake of the types and amounts of carbohydrate throughout the day, including the newer concepts of “carbohydrate counting” and the “glycemic index.” Additional attention has been paid to total calories and total fat content, plus the content of saturated fat and “trans” fat because of the importance of weight control in diabetes, and the increased risk of cardiovascular disease. **These remain the basis of diet and nutrition interventions for diabetes.**

However, information is becoming available about the special roles of **certain vitamins, minerals, phytochemicals, and some forms of dietary fat** that appear to be useful in various ways in fighting diabetes and its consequences. The nutrition tools described here do not “cure” diabetes – but in many cases they have been shown to have the capacity to **decrease the risk of diabetes developing**. Others have demonstrated that they can be useful for people who already have it by **helping with diabetic control**, or by **making diabetes hurt them less**. This includes reduction in complications like blindness, kidney failure, neuropathy, impaired circulation, and even birth defects.

**Some have been shown to be helpful in at least one of seven ways:**

- 1) Enhancing insulin sensitivity.**
- 2) Using antioxidants to minimize secondary damage from poor diabetes control caused by higher than normal production of free radicals.**
- 3) Reducing the severity of inflammation that has been shown to be associated with diabetes to decrease risk of developing diabetes (including Type I, Type II and gestational forms.)**
- 4) Helping stabilize/improve excessive fluctuations of blood sugar that can sometimes be a problem in spite of following the diet very carefully (e.g. what some people refer to as “brittle” diabetes.)**

**5) Minimizing risk of developing other threats to health and well-being, including conditions that are known to be exacerbated by diabetes.**

**6) Minimizing the risk of developing diabetes.**

**7) Discovering the emerging miscellaneous effects of a variety of phytochemicals (plant substances) on the diabetic state.**

---

Happily, all of the diet or nutrient ideas suggested here are also reported to be beneficial for most folks in terms of decreasing risks associated with one or more of a variety of chronic illnesses, including cardiovascular disease, cancer, MS, arthritis (both rheumatoid and osteo,) alzheimers, parkinsonism, depression and osteoporosis.

It is also extremely important to note that all of the suggestions are based on research reported in the legitimate peer-reviewed scientific literature. None are based on wild claims made on the internet (or elsewhere.) There is no law against fiction in America, and therefore there are quite a lot of false claims made for diet and nutrition in particular, often in the interest of selling products. Many of the “dietary supplements” sold are in fact not related to nutrition at all. Many are actually pharmaceutical products, but a loop-hole in the FDA laws allows the marketing substances in the “dietary supplement” category without having to show “safety and effectiveness” as they would if marketed as pharmaceutical products.

Understandably, the existence of so much nutritional quackery in the world has made health professionals hesitant to consider nutrition manipulations to be legitimate adjuncts to management of complex diseases like diabetes. There is a tendency to simply “throw the baby out with the bathwater.” The purpose of this paper is simply to highlight areas of research that are identifying the real “babies” to watch out for.

All the suggestions included here are generally regarded as safe at this time (I never recommend anything that comes anywhere near to “scary,”) and they are often easy to do and inexpensive. In two cases, the amount of the substance that has been shown to be beneficial is somewhat impractical to attempt when using the currently available over-the-counter products. In these cases, a prescription for the appropriate higher dosage product makes it much simpler. Although these two products and one other are also among the more expensive interventions discussed, the prescription form may be covered by insurance even though the over-the-counter form is not.

**As always, the information provided here is simply my best interpretation of the research currently available.** No claims are made that any of the large official health organizations have approved this message. It is all subject to change, which is why my papers are clearly marked with a date. And as is always the case for everything in the world “more research is ALWAYS needed,” but now there is actually quite a lot of research out there, and certainly for some things there is clearly enough evidence to initiate some safe and cheap changes. It’s at the “**won’t hurt / really might help**” level.

**I am not selling anything** – no supplements, no diet plan, no books. My only goal is to try to do some good for people. However, also as always, not all the suggestions are appropriate for every individual, so be sure to **discuss things with your personal health care provider**. **Feel free to share these materials with him/her.**

In the interests of brevity and saving trees, I will sometimes refer you to another of my papers for more complete information about a particular topic. In addition, papers with more details on this topic and others are available on the website of MeritCare Medical Center:

[www.meritcare.com](http://www.meritcare.com)

Enter my name in the search box, and then select the option that says “Cathy Breedon’s Handouts”

---

## **Part One: Enhancing insulin sensitivity**

**Magnesium and chromium** are two minerals which have specific roles in carbohydrate metabolism. **Both are frequently suboptimal in the diets of many Americans** (NHANES) for a variety of reasons and **assuring adequacy** has been shown to be critical for insulin sensitivity. For more details, references, and specific recommendations, please see “Aunt Cathy’s Guide to Nutrition: Magnesium” and “Aunt Cathy’s Guide to Nutrition: Chromium.”

**Chromium is a component of “Glucose Tolerance Factor,”** and it has roles in the metabolism of carbohydrate, protein and fat. Supplementation of chromium picolinate (a well-absorbed form) has recently been shown to be helpful in blood sugar control and weight loss among people with Type II diabetes on sulfonylurea medication. However, it may not show the same benefit among obese people on insulin whose type 2 diabetes is poorly controlled. (Diabetes Care. 2006 Mar;29(3):521-5.) In other words, it is not the only player on the team, but it is definitely one of them. It has roles in cholesterol and triglyceride metabolism, heart disease and possibly atypical depression. A review of the literature was just published on this topic: “Clinical studies on chromium picolinate supplementation in diabetes mellitus--a review.” (**Chromium references are too many to put in the text, so they follow at the end of this paper.**)

**Magnesium** has many roles in metabolizing carbohydrate, protein and fat, including being an essential cofactor in the production of ATP via the TCA Cycle. **It is specifically important for the functioning of the insulin receptors on cells**, so magnesium inadequacy is a known contributor to insulin resistance. In a very large prospective study from Harvard it was found that women with the poorest magnesium intake from food compared with the highest were 25% less likely to have developed diabetes over a 16 year period. Magnesium intake has been associated with insulin sensitivity in a threshold fashion, with a threshold at 325 mg magnesium daily. A low magnesium level in the blood (hypomagnesemia) is highly prevalent in diabetic outpatients. High plasma triglycerides, waist circumference and albuminuria are independent correlates of hypomagnesemia. The finding in the National Nutrition and Health Examination Survey (NHANES) that the majority of Americans take in less than 2/3 of the normally recommended amount suggests that improving magnesium

intake has the potential to be helpful in controlling the rapid increase in obesity and type 2 diabetes.

Interestingly, when blood sugars are high enough to **spill over into the urine**, it takes magnesium with it, resulting in an **additional loss of magnesium**. This also occurs when one uses blood pressure medications like **thiazide diuretics**. The diuretics that result in potassium loss will also result in a magnesium loss, but often this is not recognized and corrected. More will be said about magnesium later, and I also have a separate handout available with much more detail on this critical nutrient. **(Magnesium references are too many to put in the text, so they follow at the end of this paper.)**

Recently one study suggested that **vitamin K adequacy** may also have a role in both body fat and diabetes/glucose metabolism. As vitamin K inadequacy is now known to be much more common than was previously believed, the role of vitamin K in diabetes and obesity will likely begin to be evaluated more closely in the scientific community. Endocrine regulation of energy metabolism by the skeleton. Cell. 2007 Aug 10;130(3):456-69. (For more information please see my handout on the meritcare.com website for more on vitamin K: "Aunt Cathy's Guide to: Vitamin K -- New Issues in Cardiovascular Health, Osteoporosis, Cancer of the Liver and Colon, Diabetes and Varicose Veins.")

**Exercise** also has the capacity of helping to move glucose into the cells, so it is a cornerstone of prevention and also of treatment of Type II diabetes. The mineral **selenium** also appears to assist with this activity along with other functions. It is involved in the thyroid gland in setting one's resting metabolic rate, and it is also a part of a very important antioxidant in the body called glutathione peroxidase. More will be said about selenium later. **(Selenium references and Exercise references are too many to put in the text, so they follow at the end of this paper.)**

## **Part Two: Minimizing secondary damage from poor diabetes control caused by higher than normal production of free radicals**

**A greater production of free radicals is well known and documented among people with diabetes** (as well as in many other diseases with disturbed metabolism,) with the greatest production in diabetes occurring in the people with the least well-controlled blood sugar levels. Excessive production of free radicals causes serious injury to cells and tissues.

- There is evidence of **significant slowing of the development of diabetes complications by assuring a generous intake of antioxidants**. "Generous" in this context usually means substantially more than the usual RDA-type levels. The RDA guidelines are designed to meet the needs of the "healthy" population, and they have little to say about the specific needs of people with serious metabolic diseases.
- Antioxidants studied quite a lot have included **vitamins C and E, the mineral selenium** (part of the important antioxidant "glutathione peroxidase") **and a variety of carotenoid plant pigments such as lycopene in tomatoes, lutein in spinach and anthocyanin, the blue/red color in blueberries**. The latter substances have been found to be very potent antioxidants – some at 200 times the antioxidant potential of vitamin E. The ideal level

of antioxidant vitamins is unlikely to be around the RDA levels. For example, the author of a recent study concluded that “The results suggest that megadose vitamin C supplementation [1-3mg/day] may have a beneficial effect in diabetes mellitus patients on both glycemic control and antioxidant status. Thus dietary measures to increase plasma vitamin C may be an important health strategy for reducing the complications of diabetic patients.” For a review, references and recommendations for these vitamins, minerals and the carotenoid antioxidants, please see “Aunt Cathy’s Guide to Eye Health.” Other plant substances being explored for diabetes-related effects will be discussed later.

- Another unique antioxidant that has been studied extensively in diabetes is **alpha lipoic acid, (also called thioctic acid,)** a B-vitamin-like substance made in the body from the essential fatty acid linoleic acid. It also has a role in energy production, as it is required in two places in the TCA cycle. It is very benign, and the level that is generally agreed as being **most likely to bring about positive effects in diabetes research has been 600 mg/day**. A randomized, double-blind, placebo-controlled, multi-center trial showed that alpha lipoic acid at an oral dosage of 800 mg/day for 4 months significantly improved cardiac autonomic neuropathy in type 2 diabetic patients. (Treat Endocrinol. 2004;3(1):41-52. )

This is one of the three supplement substances that were described earlier as being more expensive, and it is usually available over the counter in 50 mg pills. A prescription for a higher dose pill would facilitate things. One area in which alpha lipoic acid has been most effective (and most studied) is in peripheral neuropathy research, but it now looks very interesting in several areas. (There are a great many new studies in this area with very promising results. For a list of the most recent references, see my handout Medline Search 2004-5 Studies: Diabetes x Lipoic 10/20/05 -- Short Version.)

- Another potent antioxidant with other roles in energy metabolism is **Ubiquinone – Coenzyme Q-10**. It is very safe and helpful in a number of applications, but it is also more expensive and a prescription might be helpful for this reason. CoQ10 treatment significantly improved deranged carbohydrate and lipid metabolism of experimental chemically induced diabetes in rats. The mechanism of its beneficial effect appears to be its antioxidant property. [Biol Trace Elem Res. 2006 Jan;109(1):25-34. Diabetes Res Clin Pract. 2006 Apr;72(1):100-3. Acta Diabetol. 2005 Dec;42(4):179-81. Curr Neurovasc Res. 2005 Dec;2(5):447-59.

**Research into the control of free radical production or “quenching” them after they are formed, has shown antioxidants to have promising roles in kidney health, eye circulation, lens and retinal health, circulation of blood to the extremities, wound healing, peripheral neuropathy, erectile dysfunction, the development of gestational diabetes, and birth defects. It’s time to seriously consider adding antioxidant protection to our standard protocols.** (Am J Obstet Gynecol. 2006 Feb;194(2):580-5. J Soc Gynecol Investig. 2005 Dec;12(8):549-57. Free Radic Res. 2005 Dec;39(12):1285-93. Med Hypotheses. 2006;66(1):38-44. Br J Biomed Sci. 2005;62(2):71-6.)

### **Part Three: Reducing the severity of inflammation that has been shown to be associated with diabetes**

Inflammation is now being recognized as a significant contributor to heart disease and to tissue damage in general. **Poor control of diabetes especially results in an increase in**

**inflammation.** Besides being a special problem in diabetes, this has been found to be true for many conditions that have an autoimmune component, such as rheumatoid arthritis, MS and inflammatory bowel disease. In addition to inappropriate messages sent to tissues to increase inflammation (e.g. as a result of an autoimmune condition), inflammation results from any chronic disturbances in normal fuel metabolism, such as that seen with diabetes or excessive alcohol consumption.

**Altering the ratio of omega-6 to omega-3 polyunsaturated fats appears to be an important step in achieving health for most Americans, and especially so in diabetes.** [n-3 long-chain polyunsaturated fatty acids in type 2 diabetes: a review. J Am Diet Assoc. 2005 Mar;105(3):428-40.]

Americans tend to eat about 10 g of omega-6 fat for every gram of omega-3 fat (that is, a 10-to-1 ratio.) The “Mediterranean diet” provides about a 4:1 ratio, and most healthy people would likely benefit from change in this direction. For people with diabetes, MS, and other hyperinflammatory conditions, it has been suggested that a ratio of 2:1 may provide additional benefit.

This issue has importance in minimizing complications of diabetes, risk of cancer and heart disease, and in promoting a healthy outcome of diabetic pregnancy. Other new research suggests that some individuals have a defect in producing the 20-22 carbon length fats from the shorter fat forms derived from plants, and that this may be a significant contributor to the inappropriate hyperinflammation.. For more information, references and specific recommendations, please see “Aunt Cathy’s Guide to Nutrition: Omega-3 Fats and Other Lipids.” (**Omega-3 and Omega-6 Fats references are too many to put in the text, so they follow at the end of this paper.**)

## **Part Four: Helping to stabilize excessive fluctuations of blood sugar**

Some individuals sometimes have a problem with fluctuating glucose levels in spite of following the diet, exercise and medication program very carefully (e.g. what some people used to refer to as “**brittle**” diabetes.) In the past these people were suspected of “cheating” on the diet, since we could not explain the phenomenon. Unfortunately, this accusatory response is still common in some settings.

Increasing knowledge of the effects of other **differences in types of carbohydrates consumed and factors affecting absorption** have led to some changes in our globally accepted recommendations. For example, sucrose is not looking quite so automatically bad, and the carbohydrate in the form of high fructose corn syrup is looking worse. The movement toward **whole grains/whole foods** has improved both the **nutrient:CHO ratio**, resulting in an improved magnesium and chromium intake, as described above. Increasing the intake of **soluble and insoluble fiber** in foods has positive effects, although we are quick to assume that it is the **fiber** in a “high fiber diet” that does all the good, when the diet clearly alters many other nutrition parameters as well. In any case, we no longer perceive orange juice and an intact orange to be nutritionally identical in regard to the diet for diabetes, and we now regularly use a system of **carbohydrate counting** in place of the more rigid (and much less effective) “exchange system” for meal planning.

Improved understanding of the fate of specific carbohydrate-containing foods as described above is lending some light to this puzzle, resulting in exploration of concepts such as **genetic differences** in people's metabolism, and the "**glycemic index**" of foods and meals. (For more on this, please see "Aunt Cathy's Guide to Nutrition: Issues in Childhood Obesity.")

One other important piece of the puzzle is less well known among health professionals, so I will address it here in some detail. It has been found that **relative inadequacy of carnitine exists in some people with diabetes**. [Carnitine deficiency in children and adolescents with type 1 diabetes. J Diabetes Complications. 2004 Sep-Oct;18(5):271-4 . Assessment of free L- carnitine levels in type II diabetic women with and without complications. Asia Pac J Clin Nutr. 2004;13(Suppl):S155.] This has also been found to be a factor in a number of other health conditions. Normally, one makes an adequate amount in the liver and kidney from methionine and lysine, and additional carnitine may be obtained from meat. (That's why it is called carnitine – it comes from "carne" which means meat in Latin and Spanish. Memory device: think of "chili-con-carnitine.") Individual differences in requirements, diet, genetic carnitine production, and the use of certain medications (like valproic acid for seizure control) can result in a relative inadequacy.

**Carnitine is a key component of a cellular transporter called "carnitine palmitoyl transferase" which allows fatty acids to cross the mitochondrial membrane to be used as fuel to make ATP.** Because inadequate carnitine impairs one's ability to move fatty acids into the mitochondrial membrane, symptoms can include lethargy, poor exercise endurance, poor muscle tone, cardiopathy, elevated triglyceride levels, and if applicable, breakthrough seizures. **The key feature that affects blood sugar** is the fact that there are times when we normally switch to burning primarily fat as a fuel source in order to spare glucose for the brain – sort of like having "dual heat" in your home: gas and wood. But if fat is unable to be burned because of carnitine inadequacy, a person will have to burn glucose that he/she really can't afford to burn. This results in an extremely low blood sugar, which then can trigger release of glycogen from the liver causing blood sugar to rebound up high. And if no glycogen is available to correct it, the low blood sugar can be injurious, and even life-threatening.

**Bottom line: if a person has this kind of yo-yo blood sugar with no identifiable cause, a trial on carnitine would be a good idea.** Getting a blood level is not necessarily helpful, because if it is low, then you will give some carnitine. If it is normal, blood levels do not necessarily reflect adequacy in the tissues, so if there are suggestive symptoms, one would want to do a trial anyway. The usual dose is 50-100 mg/kg/day divided into 3 doses (just because of the high osmolality,) maximum usually 3000 mg/day. To do a trial in the presence of symptoms, there is no reason not to start at the 100 mg/kg level, as the working presumption of the trial is that tissues may be depleted. Starting too low and stopping too soon may mask a true effect. When I do a trial, we monitor selected symptoms for change, and usually the trial would be continued for at least a month even in the absence of symptom relief, in order to be sure that any deficiency would have been corrected to the point of detection of symptom change.

**Carnitine is the other supplemental substance that requires a prescription for a trial and which can be expensive.** Over-the-counter products are available, but their actual

carnitine content is not assured, so at least for the trial, I would use the guaranteed product. It is also useful to consider the cost of carnitine supplementation against the cost of damage from blood sugar vacillation, and the impairment of energy production and endurance, etc., if the problem is not identified and corrected.

### **Part Five: Minimizing risk of developing other threats to health and well-being, including some conditions that are known to be exacerbated by diabetes.**

There are a number of conditions for which people with diabetes are at increased risk. These include elevated homocysteine levels, depression, cardiovascular disease (including stroke), leg cramps, and general neuropathy. Some are related to medications, such as the effect of metformin (Glucophage) on vitamin B12 level. Other common problems that can cause trouble for people with diabetes (or anyone) are these: Proton pump inhibitors for GERD may also contribute to poor absorption of vitamin B12 from food, and seriously obese people with diabetes may have had (or plan to have) gastric bypass surgery, which can also compromise vitamin B12 absorption and absorption of many nutrients. These situations will need a closer look than usual among people with diabetes.

High triglycerides are known to be a risk factor for stroke in people with diabetes especially. Three of the nutrition factors described above are associated with correcting high triglycerides: inadequate chromium intake, a high ratio of omega-6 to omega-3 fatty acids, and a relative carnitine inadequacy are all potential contributors.

The omega-6 : omega-3 ratio is also getting a lot of attention in research into cancer, MS, heart disease and arthritis, and the recommendations are all in the same direction. For more information, references and specific recommendations, please see “Aunt Cathy’s Guide to Nutrition: Omega-3 Fats and other Lipids.”

Inadequate vitamin B6 and magnesium appear to be involved in diabetic neuropathy, and also in heart disease and leg cramps. Vitamin B6, folic acid and B12 are critical for preventing elevated homocysteine, a major contributor to stroke, and possibly to alzheimers. Folic acid inadequacy is associated with cancer of the breast, prostate and colon. It is also a contributor to depression and inadequacy also makes therapeutic interventions like SSRAs work much less well. Chronic antibiotic use and alcohol abuse both impair folic acid absorption. Certain groups of people have a genetic problem that requires special attention to folic acid. For example, among some people of Irish heritage, the MTHFR gene has been discovered that contributes to higher rates of certain birth defects, depression and alcoholism. (Please see “Aunt Cathy’s Guide to Nutrition: Folic Acid” and “Aunt Cathy’s Guide to Nutrition: Vitamin B12” for details, references and recommendations.)

Some studies with vitamin B6 have used levels of 50 to 100 mg to achieve a desired effect in people with diabetic neuropathy. This is much higher than the RDA of about 2 mg, but not near a level that might be a problem. This is especially true if one is hypothesizing that their vitamin B6 requirements are in fact higher than in other people. Niacin (vitamin

B3), riboflavin (vitamin B2) and thiamine (vitamin B1) are all being studied. B vitamins are generally among the least toxic vitamins – B6 is the only one documented to cause problems with high amounts, and that was only in the most sensitive people, and never at levels below 200 mg/day. Most people who experienced any tingling in the forearms (the symptom in question) were taking over 500 mg chronically. For this reason, some people find it safe, cheaper and convenient to take a “B-100 Complex” tablet along with their general multivitamin with minerals, instead of trying to tinker with a lot of individual B vitamins. As always, check with your physician about applying any of these ideas to your own health circumstances.

As it has recently been found that serious chronic conditions like **celiac disease and hemochromatosis are much more prevalent in the general population** than has been previously thought, health professionals would do well to keep these two conditions in mind among people with diabetes. There are some links between hemochromatosis and diabetes, between hemochromatosis and celiac disease, and between diabetes and celiac disease. For example, according to a recent report, “higher iron stores (reflected by an elevated ferritin concentration and a lower ratio of transferrin receptors to ferritin) are associated with an increased risk of type 2 diabetes in healthy women independent of known diabetes risk factors.” [Body iron stores in relation to risk of type 2 diabetes in apparently healthy women. JAMA 2/2004; Coeliac disease and hereditary haemochromatosis: association and implications. Eur J Gastroenterol Hepatol. 2/2004; Hereditary haemochromatosis. QJM. 6/2004; Oxidative Stress, Beta Cell Apoptosis and Decreased Insulin Secretory Capacity in Mouse Models of Hemochromatosis. Endocrinology. 2004; Precipitation of iron overload and hereditary hemochromatosis after successful treatment of celiac disease. Am J Gastroenterol. 1/2000; Emerging concepts in celiac disease. Curr Opin Pediatr. 8/2004; Prevalence of celiac disease markers in a French cohort of children and adolescents with type 1 diabetes mellitus. Ann Endocrinol (Paris). 4/2004; Younger age at onset and sex predict celiac disease in children and adolescents with type 1 diabetes: an Italian multicenter study. Diabetes Care. 6/2004; Celiac disease associated with type 1 diabetes mellitus. Endocrinol Metab Clin North Am. 2004 Mar; others.]

**Celiac disease, an autoimmune disease that is triggered by exposure to gluten in wheat, rye and barley is genetically more common among people with Type I diabetes.** At our clinic, we screen children with Type I diabetes for hidden celiac disease with a blood test. (We also check children with Down syndrome, as they have an increased risk of both celiac disease and Type I diabetes. They have also been found to produce more free radicals than other people, and to sometimes benefit from carnitine supplementation.) In addition to the gastrointestinal problems that are the most commonly recognized symptoms of celiac disease (which can certainly affect absorption of many nutrients), it is now recognized that there are other less well recognized neurologic and dermatologic manifestations of celiac disease, and they could be complicating the recognition and treatment of neurologic and skin disorders thought to be primarily related to diabetes. Celiac is also by nature inflammatory and it also contributes to increased production of free radicals.

Both hemochromatosis and celiac disease are decidedly not rare as had been thought and both are very detrimental, so health professionals caring for people with diabetes should be alert for these problems. Hemochromatosis causes an increased risk of damage to organs like the liver, pancreas and heart, and it also is known to contribute significantly to inflammation and production of free radicals. The dietary protocol in support of regular phlebotomy can be very helpful, as I have found that many health professionals are not familiar with the many nutrition adjustments that can be of great benefit. If interested in this, please see “Aunt Cathy’s Guide to Nutrition: Hemochromatosis.”

## 6) Minimizing the risk of developing diabetes

As discussed earlier, assuring adequacy of magnesium and chromium intake may make a big difference in whether or not Type II or gestational diabetes develops in an individual. In addition, there is a major nutrition problem that is looking extremely important to correct in order to decrease the development of diabetes of any kind.

### Type I Diabetes:

There is a growing body of good evidence that **inadequacy of vitamin D in early life may be one of the triggers that brings on Type I diabetes among genetically susceptible children.** For example, in a study in Finland, children who had experienced vitamin D – deficiency rickets as infants were four times more likely to develop Type I diabetes. The data is accumulating all around the globe, and it is consistent with the well known observation that **the northern tier of the US (and places of similar latitude around the world) is known as the “Rickets Belt”, the “MS Belt” and the “Type I Diabetes Belt.”**

Latitude is the most obvious factor in vitamin D inadequacy, but many other people are at risk because of skin color, clothing that covers a person up, diet, and several other important factors. **For many reasons, vitamin D inadequacy is a very large public health problem that has not yet captured the attention of most health professionals. [I hope to be able to edit this last statement out in the near future.]** For more information, extensive references, and specific recommendations, please see “Aunt Cathy’s Guide to Nutrition: Vitamin D – It’s not just for bones any more,” and “Aunt Cathy’s Guide to Nutrition: Nutrition Issues in MS.”

### Gestational Diabetes:

Something new: Gestational diabetes appears to develop less often among women with more vitamin C in their blood. Researchers theorize that other antioxidants may have a similar effect, and other antioxidants, possibly. [Maternal plasma ascorbic Acid (vitamin C) and risk of gestational diabetes mellitus. Epidemiology. 9/2004.] Other nutrients that appear to be to be involved as well are magnesium, chromium and vitamin D. A side note: with so many young women being overweight now, and because some will actually have unrecognized Type II diabetes, it will be important to begin checking for diabetes much earlier in pregnancy (preferably before conception) in order to minimize birth defects. Women identified in a clinic as having gestational diabetes (which develops later in pregnancy) may actually have Type II diabetes all along. Please see “Aunt Cathy’s Guide to Nutrition: Top 10 Issues in Nutrition for Pregnancy” for much more on this issue.

### Type II Diabetes:

In the on-going women’s health study (Harvard), the likelihood of developing Type II diabetes among 85,000 women over a 14 year period was about 25% **less** among women who ate about 1 oz of nuts or peanuts (rich magnesium sources) 4 times a week or more compared with those who ate them rarely. The same result was seen when the risk of diabetes was evaluated relative to highest and lowest magnesium intake itself from all sources. Other studies suggest that inadequacy of vitamin D may contribute to Type II Diabetes as well as Type I, with additional roles in a wide variety of health problems such as

cancer of the colon, prostate and breast, MS, muscle/nerve pain that is often missed and diagnosed as fibromyalgia, rheumatoid and osteo arthritis, muscle weakness and falling (sarcopenia), fractures, osteoporosis, poor prenatal outcome, and much more. Assuring adequacy is easy, cheap and crucial to good health, and it is now recognized that vitamin D inadequacy is very widespread in the US and around the world.

## **7. Discovering the emerging miscellaneous effects of a variety of phytochemicals on the diabetic state.**

Around the world and in the US quite a number of plants and their various “phytochemicals” (“plant chemicals”) are being investigated for their ability to influence health in diabetes and in general. This is outside of the scope of this paper, as in general the properties being looked at are more of a pharmaceutical nature than a nutrition nature. But if you would like to get a taste of the kinds of studies being undertaken, simply log on to Public Medline ([www.nlm.nih.gov](http://www.nlm.nih.gov)) and pick the box at the right that says “for health professionals.” Then check the “Medline” box. You will see a box at the top of the screen in which you should enter search terms such as: diabetes plant or diabetes herb or diabetes phytochemical. You will find reports like this: “Coffee consumption and risk of type 2 diabetes mellitus: an 11-year prospective study of 28,812 postmenopausal women” in the journal *Arch Intern Med*. 2006 Jun 26;166(12): 1311-6. The conclusion of this study was that “Coffee intake, especially decaffeinated coffee, was inversely associated with risk of type 2 diabetes mellitus in this cohort of postmenopausal women.” That is, the coffee-drinkers had LOWER risk.

Here is just a sampling of the kinds of information available from the past few years. It’s a big topic and there has been even more research in 2007:

**2006** Effect of Tetrahydrocurcumin on Lipid Peroxidation and Lipids in Streptozotocin-Nicotinamide-Induced Diabetic Rats. *Basic Clin Pharmacol Toxicol*. 2006 Aug;99(2):122-127. Differential effects of curcumin on vasoactive factors in the diabetic rat heart. *Nutr Metab (Lond)*. 2006 Jul 18;3:27. Antioxidant effect of tetrahydrocurcumin in streptozotocin-nicotinamide induced diabetic rats. *Life Sci*. 2006 Jun 9; Effect of curcumin on protein glycosylation, lipid peroxidation, and oxygen radical generation in human red blood cells exposed to high glucose levels. *Free Radic Biol Med*. 2006 Jul 1;41(1):92-6. Curcumin attenuates thermal hyperalgesia in a diabetic mouse model of neuropathic pain. *Eur J Pharmacol*. 2006 May 1;536(3):256-61. Antidiabetic agents from medicinal plants. *Curr Med Chem*. 2006;13(10):1203-18. Hypotensive effects of hawthorn for patients with diabetes taking prescription drugs: a randomised controlled trial. *Br J Gen Pract*. 2006 Jun;56(527):437-43. The effect of the ingestion of Ginkgo biloba extract (EGb 761) on the pharmacokinetics of metformin in non-diabetic and type 2 diabetic subjects-A double blind placebo-controlled, crossover study. *Clin Nutr*. 2006 Aug;25(4):606-16. Use of herbal remedies by diabetic Hispanic women in the southwestern United States. *Phytother Res*. 2006 Apr;20(4):250-5. Puerarin protects rat pancreatic islets from damage by hydrogen peroxide. *Eur J Pharmacol*. 2006 Jan 4;529(1-3):1-7.

**2005** Plant foods in the management of diabetes mellitus: spices as beneficial antidiabetic food adjuncts. *Int J Food Sci Nutr*. 2005 Sep;56(6):399-414. Effect of supplementation of traditional medicinal plants on blood glucose in non-insulin-dependent diabetics: a pilot study. *J Med Food*. 2005 Winter;8(4):545-9. Supplementation of fenugreek leaves reduces oxidative stress in streptozotocin-induced diabetic rats. *J Med Food*. 2005 Fall;8(3):382-5. Modulatory effect of fenugreek seed mucilage & spent turmeric on intestinal & renal disaccharidases in streptozotocin induced diabetic rats. *Plant Foods Hum Nutr*. 2005 Jun;60(2):87-91. The hypoglycaemic activity of fenugreek seed extract is mediated through the stimulation of an insulin signalling pathway. *Br J Pharmacol*. 2005 Sep;146(1):41-8. Effect of fenugreek, onion and garlic on blood glucose and histopathology of pancreas of alloxan-induced diabetic rats. *Indian J Med Sci*. 2005 Feb;59(2):64-9. Antidiabetic effect of Nitobegiku, the herb *Tithonia diversifolia*, in KK-Ay diabetic mice. *Biol Pharm Bull*. 2005 Nov;28(11):2152-4. Hypoglycemic effect of ethanolic extract of *Musa sapientum* on alloxan induced diabetes mellitus in rats & its relation with antioxidant potential. *J Herb Pharmacother*. 2005;5(2):7-19. Improvement of insulin resistance by *Acanthopanax senticosus* root in fructose-rich chow-fed rats. *Clin Exp Pharmacol Physiol*. 2005 Aug;32(8):649-54. Antinociceptive, antiinflammatory & antidiabetic effects of *Leonotis leonurus* (L.) R. BR. [Lamiaceae] leaf aqueous extract in mice & rats. *Methods Find Exp Clin Pharmacol*. 2005 May;27(4):257-64. Can natural substances prevent or improve type 2 diabetes mellitus? Coffee & cinnamon for control of hyperglycemia. *MMW Fortschr Med*. 2005 Jun 23;147(25):15. Antidiabetic effect of *Cinnamomum cassia* & *Cinnamomum zeylanicum* in vivo & in vitro. *Phytother Res*. 2005 Mar;19(3):203-6. Effect of tetrahydrocurcumin on blood glucose, plasma insulin & hepatic key enzymes in streptozotocin induced diabetic rats. *J Basic Clin Physiol Pharmacol*. 2005;16(4):257-74.

**2004** Polyphenol-rich extract of fenugreek seeds protect erythrocytes from oxidative damage. *Plant Foods Hum Nutr.* 2004 Fall;59(4):143-7. What do herbalists suggest to diabetic patients in order to improve glycemic control? Evaluation of scientific evidence and potential risks. *Acta Diabetol.* 2004 Sep;41(3):91-8. Role of selected Indian plants in management of type 2 diabetes: a review. *J Altern Complement Med.* 2004 Apr;10(2):369-78. Cinnamon to control blood sugar. *MMW Fortschr Med.* 2004 Nov 25;146(48):12.

## References: Chromium

Chromium: the forgotten mineral. *Harv Mens Health Watch.* 2007 Jan;11(6):6-7; Chromium treatment has no effect in Patients with Type 2 Diabetes Mellitus in a Western Population: A Randomized, Double-Blind, Placebo-Controlled Trial. *Diabetes Care.* 2007 Feb 15; Limited evidence for effects of diet for type 2 diabetes from systematic reviews. *Eur J Clin Nutr.* 2007 Jan 24; Synergistic effects of conjugated linoleic acid and chromium picolinate improve vascular function and renal pathophysiology in the insulin-resistant JCR:LA-cp rat. *Diabetes Obes Metab.* 2007 Jan;9(1):87-95; In vitro study of elements in herbal remedies. *Biol Trace Elem Res.* 2006 Winter;114(1-3):143-50; Longitudinal hair chromium profiles of elderly subjects with normal glucose tolerance and type 2 diabetes mellitus. *Metabolism.* 2007 Jan;56(1):94-104; Relative element levels in the paired samples of scalp hair and fingernails of patients from New Delhi. *Sci Total Environ.* 2007 Jan 1;372(2-3):474-9; Chromium picolinate supplementation attenuates body weight gain and increases insulin sensitivity in subjects with type 2 diabetes: response to Martin et al. *Diabetes Care.* 2006 Dec;29(12):2764; Chromium picolinate supplementation attenuates body weight gain and increases insulin sensitivity in subjects with type 2 diabetes: response to mark. *Diabetes Care.* 2006 Dec;29(12):2764-5; Clinical studies on chromium picolinate supplementation in diabetes mellitus--a review. *Diabetes Technol Ther.* 2006 Dec;8(6):677-87; The effect of chromium picolinate and biotin supplementation on glycemic control in poorly controlled patients with type 2 diabetes mellitus: a placebo-controlled, double-blinded, randomized trial. *Diabetes Technol Ther.* 2006 Dec;8(6):636-43; Chromium picolinate intake and risk of type 2 diabetes: an evidence-based review by the United States Food and Drug Administration. *Nutr Rev.* 2006 Aug;64(8):357-63; Chromium(III) complexes of D-glucosaminic acid and their effect on decreasing blood sugar in vivo. *Arch Pharm (Weinheim).* 2006 Sep;339(9):527-30; Transcriptome of the subcutaneous adipose tissue in response to oral supplementation of type 2 Leprdb obese diabetic mice with niacin-bound chromium. *Physiol Genomics.* 2006 Nov 27;27(3):370-9; Conjugated linoleic acid and chromium lower body weight and visceral fat mass in high-fat-diet-fed mice. *Lipids.* 2006 May;41(5):437-44; Chromium picolinate supplementation attenuates body weight gain and increases insulin sensitivity in subjects with type 2 diabetes. *Diabetes Care.* 2006 Aug;29(8):1826-32; Clinical studies on chromium picolinate supplementation in diabetes mellitus--a review. *Diabetes Technol Ther.* 2006 Dec;8(6):677-87. The effect of chromium picolinate and biotin supplementation on glycemic control in poorly controlled patients with type 2 diabetes mellitus: a placebo-controlled, double-blinded, randomized trial. *Diabetes Technol Ther.* 2006 Dec;8(6):636-43. The influence of chromium chloride-containing milk to glycemic control of patients with type 2 diabetes mellitus: a randomized, double-blind, placebo-controlled trial. *Metabolism.* 2006 Jul;55(7):923-7. Chromium picolinate supplementation attenuates body weight gain and increases insulin sensitivity in subjects with type 2 diabetes. *Diabetes Care.* 2006 Aug;29(8):1826-32; Chromium picolinate positively influences the glucose transporter system via affecting cholesterol homeostasis in adipocytes cultured under hyperglycemic diabetic conditions. *Mutat Res.* 2006 Nov 7;610(1-2):93-100; The influence of chromium chloride-containing milk to glycemic control of patients with type 2 diabetes mellitus: a randomized, double-blind, placebo-controlled trial. *Metabolism.* 2006 Jul;55(7):923-7; Influence of chromium-enriched yeast on blood glucose and insulin variables, blood lipids, and markers of oxidative stress in subjects with type 2 diabetes mellitus. *Biol Trace Elem Res.* 2006 Mar;109(3):215-30; Hypoglycemic potency of novel trivalent chromium in hyperglycemic insulin-deficient rats. *J Trace Elem Med Biol.* 2006;20(1):33-9; An exploratory study into the effectiveness of a combination of traditional Chinese herbs in the management of type 2 diabetes. *Diabetes Care.* 2006 Apr;29(4):945-6; Inhibition of diabetic nephropathy in rats by an oral antidiabetic material extracted from yeast. *J Am Soc Nephrol.* 2006 Apr;17(4 Suppl 2):S127-31; Insulin-sensitizing and cholesterol-lowering effects of chromium (D-Phenylalanine)3. *J Inorg Biochem.* 2006 Jul;100(7):1187-93; Chromium treatment has no effect in patients with poorly controlled, insulin-treated type 2 diabetes in an obese Western population: a randomized, double-blind, placebo-controlled trial. *Diabetes Care.* 2006 Mar;29(3):521-5; Trivalent chromium inhibits protein glycosylation and lipid peroxidation in high glucose-treated erythrocytes. *Antioxid Redox Signal.* 2006 Jan-Feb;8(1-2):238-4; Chromium chloride inhibits TNFalpha and IL-6 secretion in isolated human blood mononuclear cells exposed to high glucose. *Horm Metab Res.* 2006 Jan;38(1):60-2; Effect of chromium picolinate on modified forced swimming test in diabetic rats: involvement of serotonergic pathways and potassium channels. *Basic Clin Pharmacol Toxicol.* 2006 Feb;98(2):155-9; Chronic CO levels have [corrected] a beneficial effect on vascular relaxation in diabetes. *Biochem Biophys Res Commun.* 2006 Feb 17;340(3):935-43; Chromium activates glucose transporter 4 trafficking and enhances insulin-stimulated glucose transport in 3T3-L1 adipocytes via a cholesterol-dependent mechanism. *Mol Endocrinol.* 2006 Apr;20(4):857-70; Mineral content of some medicinal plants used in the treatment of diabetes mellitus. *Biol Trace Elem Res.* 2005 Feb;103(2):109-15; Is chromium III a new key to effective diabetes management? *Manag Care.* 2005 Nov;14(11):78-80; Determination of fourteen trace elements in chinese traditional medicines by atomic absorption spectrometry. *Guang Pu Xue Yu Guang Pu Fen Xi.* 2005 Sep;25(9):1510-3; Mineral contents of aloe vera leaf gel and their role on streptozotocin-induced diabetic rats. *Biol Trace Elem Res.* 2005 Winter;108(1-3):185-95; Effect of chromium on the insulin resistance in patients with type II diabetes mellitus. *Folia Med (Plovdiv).* 2005;47(3-4):59-62; A double-blind, placebo-controlled, exploratory trial of chromium picolinate in atypical depression: effect on carbohydrate craving; *J Psychiatr Pract.* 2005 Sep;11(5):302-14; Effects of chronic chromium picolinate treatment in uninephrectomized rat. *Metabolism.* 2005 Sep;54(9):1243-9; Use of chromium picolinate and biotin in the management of type 2 diabetes: an economic analysis. *Dis Manag.* 2005 Aug;8(4):265-75; Supplementation with chromium picolinate recovers renal Cr concentration and improves carbohydrate metabolism and renal function in type 2 diabetic mice. *Biol Trace Elem Res.* 2005 Summer;105(1-3):229-48; Chromium supplementation shortens QTc interval duration in patients with type 2 diabetes mellitus. *Am Heart J.* 2005 Apr;149(4):632-6; Regulation of chromium on gene expression of skeletal muscles in diabetic rats. *Wei Sheng Yan Jiu.* 2005 Mar;34(2):184-7; Complementary therapies for diabetes: the case for chromium, magnesium, and antioxidants. *Arch Med Res.* 2005 May-Jun;36(3):250-7; Cellular chromium enhances activation of insulin receptor kinase. *Biochemistry.* 2005 Jun 7;44(22):8167-75; Chromium picolinate may favorably influence the vascular risk associated with smoking by combating cortisol-induced insulin resistance. *Med Hypotheses.* 2005;64(6):1220-4; Use of chromium

picolinate and biotin in the management of type 2 diabetes: an economic analysis. *Dis Manag.* **2005** Aug;8(4):265-75. *Mutat Res.* **2006** Jul 24) *Am Heart J.* **2005** Apr;149(4):632-6. *Med Hypotheses.* 2005;64(6):1220-4. *J Psychiatr Pract.* 2005 Sep;11(5):302-14. *Biol Trace Elem Res.* **2005** Summer;105(1-3):229-48.

## References: Magnesium

Effect of oral magnesium sulfate administration on blood pressure and lipid profile in streptozocin diabetic rat. *Eur J Pharmacol.* **2007** Apr 10;560(2-3):201-5. *Epub* **2007** Jan 19; Magnesium metabolism in type 2 diabetes mellitus, metabolic syndrome and insulin resistance. *Arch Biochem Biophys.* **2007** Feb 1;458(1):40-7. The effect of magnesium deficit on serum immunoglobulin concentrations in type 1 diabetes mellitus. *Rom J Intern Med.* **2006**;44(1):61-7; Serum magnesium and type-2 diabetes in African Americans and Hispanics: a New York cohort. *J Am Coll Nutr.* **2006** Dec;25(6):509-13. Magnesium intake is related to improved insulin homeostasis in the framingham offspring cohort. *J Am Coll Nutr.* **2006** Dec;25(6):486-92; Increased magnesium intake prevents hyperlipidemia and insulin resistance and reduces lipid peroxidation in fructose-fed rats *Pathophysiology.* **2006** Dec 20; Relationship between serum magnesium values, lipids and anthropometric risk factors. *Atherosclerosis.* **2006** Dec 7; Dietary magnesium intake and risk of incident hypertension among middle-aged and older US women in a 10-year follow-up study. *Am J Cardiol.* **2006** Dec 15;98(12):1616-21; Other relevant components of nuts: phytosterols, folate and minerals. *Br J Nutr.* **2006** Nov;96 Suppl 2:S36-44; Intracellular magnesium in elderly patients with heart failure: effects of diabetes and renal dysfunction. *J Trace Elem Med Biol.* **2006**;20(4):221-6; Dietary magnesium and fiber intakes and inflammatory and metabolic indicators in middle-aged subjects from a population-based cohort. *Am J Clin Nutr.* **2006** Nov;84(5):1062-9; Dietary calcium and magnesium, major food sources, and risk of type 2 diabetes in U.S. black women. *Diabetes Care.* **2006** Oct;29(10):2238-43; Dietary fiber intake, dietary glycemic load, and the risk for gestational diabetes mellitus. *Diabetes Care.* **2006** Oct;29(10):2223-30; Effects of oral magnesium supplementation on glycaemic control in Type 2 diabetes: a meta-analysis of randomized double-blind controlled trials. *Diabet Med.* **2006** Oct;23(10):1050-6. Dairy, magnesium, and calcium intake in relation to insulin sensitivity: approaches to modeling a dose-dependent association. *Am J Epidemiol.* **2006** Sep 1;164(5):449-58. The nerve-heart connection in the pro-oxidant response to Mg-deficiency. *Heart Fail Rev.* **2006** Mar;11(1):35-44; A prospective study of dairy intake and the risk of type 2 diabetes in women. *Diabetes Care.* **2006** Jul;29(7):1579-84; Serum ionized magnesium levels in relation to metabolic syndrome in type 2 diabetic patients. *J Am Coll Nutr.* **2006** Jun;25(3):210-5; Magnesium and diabetes mellitus: their relation. *Clin Nutr.* **2006** Aug;25(4):554-62. Magnesium intake and incidence of metabolic syndrome among young adults. *Circulation.* **2006** Apr 4;113(13):1675-82; Dietary intake among youth with diabetes: the SEARCH for Diabetes in Youth Study. *J Am Diet Assoc.* **2006** May;106(5):689-97.; *Am J Epidemiol.* 2006 Jul 21; *J Am Coll Nutr.* 2006 Jun;25(3):210-5. *Clin Nutr.* 2006 Aug;25(4):554-62. *Arch Biochem Biophys.* **2006** Jun 12; Magnesium urinary excretion in diabetic adolescents. *Acta Medica (Hradec Kralove).* **2005**;48(3-4):157-61

## References: Selenium

Differential effects of selenium compounds on glucose synthesis in rabbit kidney-cortex tubules and hepatocytes. In vitro and in vivo studies. *J Inorg Biochem.* **2007** Mar;101(3):493-505; Maternal-fetal status of copper, iron, molybdenum, selenium, and zinc in obese pregnant women in late gestation. *Biol Trace Elem Res.* **2006** Nov;113 (2):113-23; Antioxidant supplementation does not affect fasting plasma glucose in the Supplementation with Antioxidant Vitamins and Minerals (SU.VI.MAX) study in France: association with dietary intake and plasma concentrations. *Am J Clin Nutr.* **2006** Aug;84(2):395-9; Selenium and glutathione peroxidase enzyme levels in diabetic patients with early spontaneous abortions. *Akush Ginekol (Sofia).* **2006**;45(5):3-9; Trace element status in Saudi patients with established atherosclerosis. *J Trace Elem Med Biol.* **2006**;20(2):105-14; Maternal-fetal transport and disposition of copper, iron, molybdenum, selenium and zinc in experimentally induced diabetic rats. *J Matern Fetal Neonatal Med.* **2006** Jan;19(1):57-64; Influence of selenium (antioxidant) on gliclazide induced hypoglycaemia/anti hyperglycaemia in normal/alloxan-induced diabetic rats. *Mol Cell Biochem.* **2006** Feb;283(1-2):123-7; Compendium of the antidiabetic effects of supranutritional selenate doses. In vivo and in vitro investigations with type II diabetic db/db mice. *J Nutr Biochem.* **2006** Aug;17(8):548-60; The determinants of endothelial dysfunction in CKD: oxidative stress and asymmetric dimethylarginine. *Am J Kidney Dis.* **2006** Jan;47(1):42-50; Toenail cerium levels and risk of a first acute myocardial infarction: the EURAMIC and heavy metals study. *Chemosphere.* **2006** Jun;64(1):112-20; Effect of selenite treatment on ultrastructural changes in experimental diabetic rat bones. *Biol Trace Elem Res.* **2005** Nov;107(2):167-79; Selenium prevents diabetes-induced alterations in [Zn<sup>2+</sup>]<sub>i</sub> and metallothionein level of rat heart via restoration of cell redox cycle. *Am J Physiol Heart Circ Physiol.* **2006** Mar;290(3):H1071-80; Effect of the joint administration of selenium and vitamin E in combination with regular aerobic exercise on markers of lipid peroxidation and glutathione peroxidase in diabetic rats. *Int J Sport Nutr Exerc Metab.* **2005** Jun;15(3):266-78; Toenail selenium and cardiovascular disease in men with diabetes. *J Am Coll Nutr.* **2005** Aug;24(4):250-6; Selenium treatment protects diabetes-induced biochemical and ultrastructural alterations in liver tissue. *Biol Trace Elem Res.* **2005** Summer; 105(1-3):135-50; Glucooxidative stress and spontaneous abortion in pregnant women with diabetes mellitus type. *Akush Ginekol (Sofia).* **2005**;44(4):3-10; Embryopathy in experimental diabetic gestation: assessment of oxidative stress and antioxidant defence. *Br J Biomed Sci.* **2005**;62(2):71-6; Influence of alloxan-induced diabetes and selenite treatment on blood glucose and glutathione levels in mice. *J Trace Elem Med Biol.* **2005**;18(3):261-7; Effect of alpha-tocopherol on the metabolic control and oxidative stress in female type 2 diabetics. *Biomed Pharmacother.* **2005** Jul;59(6):290-5; Diabetes mellitus worsens antioxidant status in patients with chronic pancreatitis. *Am J Clin Nutr.* **2005** May;81(5):1117-25; Nutritional, dietary and postprandial oxidative stress. *J Nutr.* **2005** May;135(5):969-72. Effects of a supranutritional dose of selenate compared with selenite on insulin sensitivity in type II diabetic db/db mice. *J Anim Physiol Anim Nutr (Berl).* **2005** Apr-Jun;89(3-6):94-104; Interpretation of relevance of sodium-calcium exchange in action potential of diabetic rat heart by mathematical model. *Mol Cell Biochem.* **2005** Jan;269(1-2):121-9; Beneficial effects of selenium on some enzymes of diabetic rat heart. *Biol Trace Elem Res.* **2005** Mar;103(3):207-16; Selenium and endocrine systems. *J Endocrinol.* **2005** Mar;184(3):455-65; The role of antioxidant micronutrients in the prevention of diabetic complications. *Treat Endocrinol.*

2004;3(1):41-52; Gestational hyperglycemia, zinc, selenium, and antioxidant vitamins. *Nutrition*. 2005 Feb;21(2):186-91; What do herbalists suggest to diabetic patients in order to improve glycemic control? Evaluation of scientific evidence and potential risks. *Acta Diabetol*. 2004 Sep;41(3):91-8; The putative therapeutic value of high-dose selenium in proliferative retinopathies may reflect down-regulation of VEGF production by the hypoxic retina. *Med Hypotheses*. 2005;64(1):159-6; Maternal-fetal status of copper, iron, molybdenum, selenium and zinc in insulin-dependent diabetic pregnancies. *Arch Gynecol Obstet*. 2005 Mar;271(3):212-7; Effects of sodium selenate treatment on altered responses of left and right atria from streptozotocin-induced diabetic rats. *J Cardiovasc Pharmacol*. 2004 Jul;44(1):9-15; Effects of selenium on altered mechanical and electrical cardiac activities of diabetic rat. *Arch Biochem Biophys*. 2004 Jun 1;426(1):83-90; Effects of zinc and selenium on the disorders of blood glucose and lipid metabolism and its molecular mechanism in diabetic rats. *Wei Sheng Yan Jiu*. 2004 Jan;33(1):70-3; Maternal-fetal status of copper, iron, molybdenum, selenium and zinc in patients with gestational diabetes. *J Matern Fetal Neonatal Med*. 2004 Jul;16(1):15-21; Plasma selenium decrease during pregnancy is associated with glucose intolerance. *Biol Trace Elem Res*. 2004 Jul;100(1):19-29. Selenium supplementation decreases nuclear factor-kappa B activity in peripheral blood mononuclear cells from type 2 diabetic patients. *Eur J Clin Invest*. 2004 Jul;34(7):475-81. New experimental observation on the relationship of selenium and diabetes mellitus. *Biol Trace Elem Res*. 2004 Summer;99(1-3):241-53; Development of insulin resistance and obesity in mice overexpressing cellular glutathione peroxidase. *Proc Natl Acad Sci U S A*. 2004 Jun 15;101(24):8852-7; Organoselenium compounds as potential therapeutic and chemopreventive agents: a review. *Curr Med Chem*. 2004 Jun;11(12):1657-69; The chemical form of selenium affects insulinomimetic properties of the trace element: investigations in type II diabetic dbdb mice. *J Nutr Biochem*. 2003 Nov;14(11):637-47; Activities of xanthine oxidoreductase and antioxidant enzymes in different tissues of diabetic rats. *J Lab Clin Med*. 2003 Sep;142(3):172-7; Protective effects of antioxidant micronutrients (vitamin E, zinc and selenium) in type 2 diabetes mellitus. *Clin Chem Lab Med*. 2003 Aug;41(8):995-8; Effect of sodium selenite treatment on platelet aggregation of streptozotocin-induced diabetic rats. *Thromb Res*. 2003;111(6):363-7.

## References: Exercise

Effects of aerobic exercise on microalbuminuria and enzymuria in type 2 diabetic patients. *Ren Fail*. 2007;29(2):199-205; Diabetes mellitus, part 1: physiology and complications. *Br J Nurs*. 2007 Feb 8-21;16(3):184-8; Mexican Americans with type 2 diabetes: perspectives on definitions, motivators, and programs of physical activity. *Prev Chronic Dis*. 2007 Apr;4(2):A24. Epub 2007 Mar 15; Variation in GYS1 Interacts with Exercise and Gender to Predict Cardiovascular Mortality. *PLoS ONE*. 2007 Mar 14;2:e285; Exercise capacity and body mass as predictors of mortality among male veterans with type 2 diabetes. *Diabetes Care*. 2007 Mar 10; Correlates of Physical Activity in Pregnancy among Latina Women. *Matern Child Health J*. 2007 Mar 8; Skeletal muscle lipid deposition and insulin resistance: effect of dietary fatty acids and exercise. *Am J Clin Nutr*. 2007 Mar;85(3):662-77; Metabolic effects of interventions to increase exercise in adults with type 2 diabetes. *Diabetologia*. 2007 Mar 7; Triple jeopardy: nocturnal hypoglycemia after exercise in the young with diabetes. *J Clin Endocrinol Metab*. 2007 Mar;92(3):815-6; Artificial Selection for High Capacity Endurance Running is Protective Against High Fat Diet-Induced Insulin Resistance. *Am J Physiol Endocrinol Metab*. 2007 Mar 6; Effect of intermittent high-intensity compared with continuous moderate exercise on glucose production and utilization in individuals with type 1 diabetes. *Am J Physiol Endocrinol Metab*. 2007 Mar;292(3):E865-70; Effect of moderate aerobic exercise on sympatho-vagal balance in Type 2 diabetic patients. *Diabet Med*. 2007 Feb 28; Physical activity in prevention and treatment of the metabolic syndrome. *Appl Physiol Nutr Metab*. 2007 Feb;32(1):76-88; Impact of the short-term, intense exercise on postprandial glycemia in type 2 diabetic patients treated with gliclazide. *J Diabetes Complications*. 2007 Mar-Apr;21(2):101-7; Type 2 diabetes and cardiovascular disease: reducing the risk. *J Manag Care Pharm*. 2007 Mar;13(2 Supp A):2-15; Therapy insight: peripheral arterial disease and diabetes--from pathogenesis to treatment guidelines. *Nat Clin Pract Cardiovasc Med*. 2007 Mar;4(3):151-62. Effect of Acute Exercise on AMPK Signaling in Skeletal Muscle of Subjects With Type 2 Diabetes: A Time-Course and Dose-Response Study. *Diabetes*. 2007 Mar;56(3):836-48; Genetic Variations in PPAR $\delta$  and PPAR $\gamma$ 1A determine Mitochondrial Function and Change in Aerobic Physical Fitness and Insulin Sensitivity during Lifestyle Intervention. *J Clin Endocrinol Metab*. 2007 Feb 27; Exercise reduces resistin and inflammatory cytokines in patients with type 2 diabetes. *Diabetes Care*. 2007 Mar;30(3):719-21; Habitual physical activity is associated with intrahepatic fat content in humans. *Diabetes Care*. 2007 Mar;30(3):683-8; High-intensity training improves plasma glucose and acid-base regulation during intermittent maximal exercise in type 1 diabetes. *Diabetes Care*. 2007 Feb 26; Identification of Distinct Self-Management Styles of Adolescents with Type 1 Diabetes. *Diabetes Care*. 2007 Feb 23; Exercise Interventions for Diabetes Control: Do We Really Know That Strength Training Is Better Than Endurance Training? *Arch Phys Med Rehabil*. 2007 Mar;88(3):397; Renal functional changes in relation to hemodynamic parameters during exercise test in normoalbuminuric insulin-dependent children. *Acta Paediatr*. 2007 Feb 13; The role of exercise in patients with type 2 diabetes. *Am Fam Physician*. 2007 Feb 1;75(3):335-6; Walking decreased risk of cardiovascular disease mortality in older adults with diabetes. *J Clin Epidemiol*. 2007 Mar;60(3):309-17; Diets and discipline: the narratives of practice of university students with type 1 diabetes. *Social Health Illn*. 2007 Jan;29(1):136-53; Physical activity and metabolic risk in individuals with a family history of type 2 diabetes. *Diabetes Care*. 2007 Feb;30(2):337-42; Physical activity in U.S. Adults with diabetes and at risk for developing diabetes, 2003. *Diabetes Care*. 2007 Feb;30(2):203-9; Exercise for prevention and treatment of cardiovascular disease, type 2 diabetes, and metabolic syndrome. *Curr Diab Rep*. 2007 Feb;7(1):14-9; Substrate oxidation during exercise: type 2 diabetes is associated with a decrease in lipid oxidation and an earlier shift towards carbohydrate utilization. *Diabetes Metab*. 2006 Dec;32(6):604-10; A physical activity programme and its effects on insulin resistance and oxidative defense in obese male patients with type 2 diabetes mellitus. *Diabetes Metab*. 2006 Dec;32(6):583-90.

## References: Omega-3 and Omega-6 Fats

Postprandial effect of n-3 polyunsaturated fatty acids on apolipoprotein B-containing lipoproteins and vascular reactivity in type 2 diabetes. *Am J Clin Nutr*. 2007 Feb;85(2):369-76; Dietary omega-3 fatty acids for women. *Biomed Pharmacother*. 2007 Jan 2; Blood omega-3 and trans fatty acids in middle-aged acute coronary syndrome patients. *Am J Cardiol*. 2007 Jan 15;99(2):154-8; Meta-analysis of the effects of n-3 polyunsaturated fatty acids on haematological and thrombogenic factors in type 2 diabetes. *Diabetologia*.

2007 Feb;50(2):250-8; Nutrition and inflammatory load. *Ann N Y Acad Sci.* 2006 Nov;1083:214-38. Omega-3 fatty acids and mood disorders. *Am J Psychiatry.* 2006 Nov;163(11): 2018; Brain phospholipid and triglyceride fatty acid content and pattern in Type 1 and Type 2 diabetic rats. *Neurosci Lett.* 2006 Nov 27;409 (1):75-9; Lipometabolic disorder--cholesterol and triglycerides. *Clin Res Cardiol.* 2006;95 Suppl 6:V117-22; Fish oil and mental health: the role of n-3 long-chain polyunsaturated fatty acids in cognitive development and neurological disorders. *Int Clin Psychopharmacol.* 2006 Nov;21(6): 319-36; Cardiac cell death in early diabetes and its modulation by dietary fatty acids. *Biochim Biophys Acta.* 2006 Oct;1761(10):1148-62; Differential effects of n-3 polyunsaturated fatty acids on metabolic control and vascular reactivity in the type 2 diabetic ob/ob mouse. *Metabolism.* 2006 Oct;55(10):1365-74; Effects of n-3 fatty acids in subjects with type 2 diabetes: reduction of insulin sensitivity and time-dependent alteration from carbohydrate to fat oxidation. *Am J Clin Nutr.* 2006 Sep;84(3):540-50; N-3 fatty acid supplementation decreases plasma homocysteine in diabetic dyslipidemia treated with statin-fibrate combination. *J Nutr Biochem.* 2006 Jun;17(6):379-84; Gamma linolenic acid: an antiinflammatory omega-6 fatty acid. *Curr Pharm Biotechnol.* 2006 Dec;7(6):531-4; Depression and cardiovascular mortality: a role for n-3 fatty acids? *Am J Clin Nutr.* 2006 Dec;84(6):1513-7; The metabolic syndrome, omega-3 fatty acids and inflammatory processes in relation to schizophrenia. *Prostaglandins Leukot Essent Fatty Acids.* 2006 Oct-Nov;75(4-5):323-7; Dietary fish and n-3 fatty acid intake and cardiac electrocardiographic parameters in humans. *J Am Coll Cardiol.* 2006 Aug 1;48(3):478-84; Effect of n-3 fatty acids on metabolism of apoB100-containing lipoprotein in type 2 diabetic subjects. *Br J Nutr.* 2006 Jul;96(1):100-6; n-3 fatty acids and gene expression. *Am J Clin Nutr.* 2006 Jun;83(6 Suppl):1520S-1525S; n-3 fatty acids and the metabolic syndrome. *Am J Clin Nutr.* 2006 Jun;83(6 Suppl):1499S-1504S; Hypertriglyceridemia: management of atherogenic dyslipidemia. *J Fam Pract.* 2006 Jul;55(7):S1-8; N-3 fatty acids modulate Th1 and Th2 dichotomy in diabetic pregnancy and macrosomia. *J Autoimmun.* 2006 Jun;26(4):268-77; Hypertriglyceridemia. *J Am Board Fam Med.* 2006 May-Jun;19(3): 310-6; Effect of n-3 and n-6 polyunsaturated fatty acids on lymphocyte proliferation, interleukin production and phospholipid fatty acids composition in type 2 diabetic and healthy subjects in Jordan people. *Prostaglandins Leukot Essent Fatty Acids.* 2006 Jun;74(6):347-56; The relationship between tissue RBC n-3 fatty acids and pulse wave velocity. *In Vivo.* 2006 Mar-Apr;20(2):307-10; Association between dietary fat and anti-oxidant status of Tunisian type 2 diabetic patients. *Prostaglandins Leukot Essent Fatty Acids.* 2006 May;74(5):323-9; Intake of fish and long-chain n-3 fatty acids and the risk of coronary heart mortality in men and women. *Br J Nutr.* 2006 Apr;95(4):824-9; Sustained enrichment of liver phospholipids and triglycerides in eicosapentaenoate after a bolus intravenous injection of a medium-chain triglycerides: fish oil emulsion to streptozotocin (Type 1) and Goto-Kakizaki (Type 2) diabetic rats. *Int J Mol Med.* 2006 Apr;17(4):643-7; Nutrient intake of women with and without gestational diabetes with a specific focus on fatty acids. *Nutrition.* 2006 Mar;22(3):230-6; A distinctive fatty acid profile in circulating lipids of Korean gestational diabetics: a pilot study. *Diabetes Res Clin Pract.* 2006 Aug;73(2):178-83; N-3 fatty acids modulate antioxidant status in diabetic rats and their macroscopic offspring. *Int J Obes (Lond).* 2006 May;30(5):739-50; Differential effect of saturated, monounsaturated, and polyunsaturated fatty acids on alloxan-induced diabetes mellitus. *Prostaglandins Leukot Essent Fatty Acids.* 2006 Mar;74(3):199-213; Effect of n-3 polyunsaturated fatty acids on plasma lipid, LDL lipoperoxidation, homocysteine and inflammation indicators in diabetic dyslipidemia treated with statin + fibrate combination. *Cas Lek Cesk.* 2005;144(11):737-41; Dietary fats, fatty acids and insulin resistance: short review of a multi-faceted connection. *Med Sci Monit.* 2005 Dec;11(12):RA359-67; A successful diabetes prevention study in Eskimos: the Alaska Siberia project. *Int J Circumpolar Health.* 2005 Sep;64(4):409-24; Omega-3 fatty acids improve glucose tolerance and components of the metabolic syndrome in Alaskan Eskimos: the Alaska Siberia project. *Int J Circumpolar Health.* 2005 Sep;64(4):396-408; Fat food for a bad mood. Could we treat and prevent depression in Type 2 diabetes by means of omega-3 polyunsaturated fatty acids? A review of the evidence. *Diabet Med.* 2005 Nov;22 (11):1465-75. Mediation of cognitive function by high fat diet following stress and inflammation. *Nutr Neurosci.* 2005 Oct-Dec;8(5-6):309-15; Nutrition in prediabetes. *J Indian Med Assoc.* 2005 Nov;103(11):596, 598-9; C-reactive proteins and chronic disease: what role does nutrition play? *Nutr Clin Pract.* 2003 Jun;18(3):227-33; Molecular mechanisms of action and health benefits of polyunsaturated fatty acids] *Rev Invest Clin.* 2005 May-Jun;57(3):457-72; Effects of omega-3 fatty acids on eye health. *Evid Rep Technol Assess (Summ).* 2005 Jul;(117):1-6; Extending the cardiovascular benefits of omega-3 Fatty acids. *Curr Atheroscler Rep.* 2005 Sep;7(5):375-80; The fats of life: type 1 diabetes mellitus, athero-sclerosis, & omega-3 long-chain polyunsaturated fatty acids. *Treat Endocrinol.* 2005;4(4):261; The impact of age, body mass index, & fish intake on the EPA & DHA content of human erythrocytes. *Lipids.* 2005 Apr;40(4):343-7; Structured dietary advice incorporating walnuts achieves optimal fat and energy balance in patients with type 2 diabetes mellitus. *J Am Diet Assoc.* 2005 Jul;105(7):1087-96; Fetal erythrocyte membrane lipids modification: preliminary observation of an early sign of compromised insulin sensitivity in offspring of gestational diabetic women. *Diabet Med.* 2005 Jul;22(7):914-20; Omega-3 fatty acids in the treatment of psychiatric disorders. *Drugs.* 2005;65(8):1051-9; Ratio of n-6 to n-3 fatty acids & bone mineral density in older adults: the Rancho Bernardo Study. *Am J Clin Nutr.* 2005 Apr;81(4):934-8; Long-chain omega 6 polyunsaturated fatty acids in erythrocyte phospholipids are associated with insulin resistance in non-obese type 2 diabetics. *Clin Chim Acta.* 2005 Apr;354(1-2):195-9; n-3 long-chain polyunsaturated fatty acids in type 2 diabetes: a review. *J Am Diet Assoc.* 2005 Mar;105(3):428-40; A few more thoughts about fish and fish oil. *J Am Diet Assoc.* 2005 Mar;105(3):350-1; Red wine consumption improves insulin resistance but not endothelial function in type 2 diabetic patients. *Metabolism.* 2005 Mar;54(3):306-13; Dietary linolenic acid is associated with a lower prevalence of hypertension in the NHLBI Family Heart Study. *Hypertension.* 2005 Mar;45(3):368-73; Omega-3 fatty acids: molecular approaches to optimal biological outcomes. *Curr Opin Lipidol.* 2005 Feb;16(1):11-8; Acute effects of monounsaturated fatty acids with and without omega-3 fatty acids on vascular reactivity in individuals with type 2 diabetes. *Diabetologia.* 2005 Jan;48(1):113-22; The role of omega-3 long-chain polyunsaturated fatty acids in health and disease of the retina. *Prog Retin Eye Res.* 2005 Jan;24(1):87-138; Including walnuts in a low-fat/modified-fat diet improves HDL cholesterol-to-total cholesterol ratios in patients with type 2 diabetes. *Diabetes Care.* 2004 Dec;27(12):2777-83.